Counter Intelligence PLUS | 2025

The training guide for pharmacy assistants



Phenergan 10mg tablets Promethazine hydrochloride Oral use

Phenergan 10mg Tablets

Uses

Treatment of allergic conditions such as hayfever or rashes such as nettle rash or hives. Also for the treatment of short term adult insomnia, for short-term use as a paediatric sedative and to treat or stop nausea or vomiting such as travel sickness.

Dosage

For allergies: Adults, the elderly and children over 10 years: one tablet twice a day. This may be increased to a maximum of two tablets three times a day. Children 5-10 years: a single dose of either one or two tablets given at night or one tablet given twice a day. No more than two tablets a day. Children 2-5 years: Phenergan Elixir recommended. Children under 2 years: not recommended. As an anti-emetic: Adults, the elderly and children over 10 years: two tablets the night before the journey. Repeat after 6-8 hours as required. Children 5-10 years: one tablet the night before the journey. Repeat after 6-8 hours as required. Children 2-5 years: Phenergan Elixir recommended. Children under 2 years: not recommended. For short term treatment of insomnia in adults and for short term use as a paediatric sedative: Children over 10 years and adults (including the elderly): 2-5 tablets (20mg

10 years and adults (including the elderly): 2-5 tablets (20mg to 50mg) as a single dose at nighttime. Children 5-10 years: 2 tablets (20mg) given as a single dose at nighttime. Children 2-5 years: Phenergan Elixir should be given in this age group.

Format and Ingredients

Film coated tablet containing: Promethazine hydrochloride 10mg,

Category:

Pack size: 56



Communications International Group Ltd 2025, Linen Hall, 162-168 Regent Street, London W1B 5TB