

Counter Intelligence PLUS | 2025



The training guide for pharmacy assistants



Solpadeine Max Soluble Tablets

Uses

For the short term treatment of acute moderate pain which is not relieved by paracetamol or ibuprofen alone. Recommended for the relief of headache, migraine, dental pain, period pain, strains and sprains, backache, arthritic and rheumatic pain and sciatica.

Dosage

Over 18 years: 2 tablets dissolved in at least half a tumbler of water every 4-6 hours as required but not more frequently than every 4 hours. No more than 8 tablets in 24 hours. 16-18 years: 1-2 tablets dissolved in at least half a tumbler of water every 6 hours as required but not more frequently than every 6 hours. No more than 8 tablets in 24 hours. 12-15 years: 1 tablet dissolved in at least half a tumbler of water every 6 hours as required but not more frequently than every 6 hours. No more than 4 tablets in 24 hours. The duration of treatment should be limited to 3 days. 12 years and under: not to be given.

Format and Ingredients

Effervescent tablet containing: Paracetamol 500mg, Codeine Phosphate Hemihydrate 12.8mg, Caffeine 30mg, Analgesics Pack size: 32

Manufacturer: RRP: £8.59 PERRIGO

Legal Status: (P)

A division of GROUPE EUROCOM



