



Counter Intelligence PLUS | 2025

The training guide for pharmacy assistants



ANBESOL®

teething gel

Lidocaine Hydrochloride,
Chlorocresol,
Cetylpyridinium Chloride.

Effective soothing relief from: • teething pain • mouth ulcers • denture irritation



FOR BABIES
AND ADULTS

SUGAR FREE

Anbesol Teething Gel

Uses

Adults and the elderly: temporary relief of pain caused by recurrent mouth ulcers and denture irritation. Children over 5 months: for relief of pain and discomfort associated with teething where non-pharmacological options have failed.

Dosage

Adults and the elderly: apply a small amount to affected area with a clean fingertip. Two immediate applications normally sufficient. Not to be used more frequently than every 3 hours. Babies and children over 5 months: apply a small amount to affected area with a clean fingertip. The dose may be repeated if necessary after 3 hours, up to a maximum of 6 doses in 24 hours. Treatment should be stopped once symptoms have resolved. Not to be used for more than 7 days.

Format and Ingredients

Gel containing:

Lidocaine Hydrochloride 1.0mg, Chlorocresol 0.1mg,
Cetylpyridinium Chloride 0.02mg,

Category:

Analgesics, Infants and Babies

Manufacturer:

ALLIANCE PHARMACEUTICALS
LTD

Pack size: 10g

RRP: £2.39

Legal Status: (P)